

## Original Research Article

# Study of serum lactate dehydrogenase level as diagnostic and prognostic indicator of megaloblastic anemia

Neelendra Chakravarty<sup>1</sup>, D. Santhikiran<sup>1\*</sup>, Anupam Brahma<sup>1</sup>, U. R. Singh<sup>3</sup>, P. C. Kol<sup>3</sup>,  
Sukanta Sen<sup>2</sup>

<sup>1</sup>Department of Pathology, <sup>2</sup>Department of Pharmacology, ICARE Institute of Medical Sciences and Research, Haldia, West Bengal, India

<sup>3</sup>Department of Pathology, Shyam Shah Medical College, Rewa, Madhya Pradesh, India

**Received:** 21 April 2019

**Accepted:** 30 May 2019

### \*Correspondence:

Dr. D. Santhikiran,

E-mail: [dr.dsanthikiran@gmail.com](mailto:dr.dsanthikiran@gmail.com)

**Copyright:** © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## ABSTRACT

**Background:** Megaloblastic anaemia is the hematologic manifestation of faulty proliferation of blood cell precursors. The present study was done to facilitate the diagnosis prior to performing any bone marrow aspirate by estimation of the value of serum LDH in the diagnosis of megaloblastic anaemia.

**Methods:** The cases were selected from patients attended the OPD and admitted in Sanjay Gandhi Memorial Hospital & Gandhi Memorial Hospital, Shyam Shah Medical College, Rewa, Madhya Pradesh. Following investigations were then done to classify anaemia and to establish the diagnosis of megaloblastic anaemia like Haemoglobin estimation by cyanmethaemoglobin method, PCV, RBC count and absolute values, general blood picture, reticulocyte count, bone marrow examination and serum LDH estimation before and after treatment.

**Results:** Of the 100 cases, 50 cases (50%) of the cases were microcytic hypochromic anaemia. 15 cases (15%) were normocytic normochromic anaemia; 35 cases (35%) were macrocytic anaemia on the basis of general blood picture and absolute values. The incidence of megaloblastic anaemia in Indian adults was 20%. Maximum number of cases (90%) of the cases had serum LDH level of more than 1000 U/L. Range of serum LDH level was 520 U/L to 4520 U/L. Thus, there was 2 to 20-fold of highest reference value (240 U/L at 37°C) rise in serum LDH level in megaloblastic anaemia.

**Conclusions:** Megaloblastic anaemia is not uncommon in Indian adults and serum LDH levels provide an important means of diagnosis. It is a non-invasive procedure, safe, and does not require any expertise.

**Keywords:** Central India, Dehydrogenase, Hemoglobin, Megaloblastic anaemia, Serum lactate

## INTRODUCTION

Anemia is present when the hemoglobin level in the blood is below the lower extreme of the normal range for the age and sex of the individual. Megaloblastic anemia is the hematologic manifestation of faulty proliferation of blood cell precursors.<sup>1-3</sup> More than 95% of megaloblastic macrocytic anemia is the result of a deficiency of vitamin

B12 and folates.<sup>4</sup> In India, according to a report of the meeting "Consultation on nutritional anemia", 15-25% of adult males, 50-80% of adult females 76% preschool children, 50% school children were found to be suffering from nutritional anemia.<sup>5</sup>

Among the nutritional requirement for red cell production, Iron, Vitamin B12 and folic acid are most important. Their

deficiency leads to nutritional anemia. They are the morphologically similar group of pancytopenic disorders having in common a selective reduction in the rate of DNA synthesis.<sup>6</sup> The macrocytic anemias are characterized by an increase in the average size of erythrocyte. Typically, all dimensions of the cell are increased. The amount of hemoglobin in each cell is increased in proportion to the size. Macrocytosis is best detected by determination of mean corpuscular volume. Values exceeding 100 fl are generally considered to establish the presence of macrocytosis. Mean corpuscular hemoglobin value is also increased but the mean corpuscular hemoglobin concentration remains within normal limits.<sup>7</sup> Although most common than any anemia is due to iron deficiency, they are a significant cause of ill health in many parts of the world.<sup>8</sup>

Definite diagnosis of megaloblastic anemia is made by bone marrow examination and demonstration of characteristic megaloblasts. They have large size and delicate sieve-like nuclear chromatin. An unusually large number of mitotic figures are found among the erythroid cells. Elevated serum LDH are observed in a variety of conditions. The highest values (two to forty-fold elevations) are seen in patients with megaloblastic anemia. Intramedullary destruction of immature megaloblastic cells has been suggested as the cause of increased release of enzyme from the bone marrow. It is not only the increased intramedullary turnover of megaloblastic cells but also a higher LDH content of these cells that is responsible for high LDH plasma levels.<sup>9,10</sup>

Appropriate treatment of the megaloblastic anemia with either vitamin B12 or folic acid produces a dramatic reduction of serum LDH activity. Which proceeds increase the reticulocyte count by several days and showed that the LDH present in pernicious anemia serum had a marked preponderance of the fast-moving isoenzymes LDH-1 and LDH-2, with LDH-1 activity greater than LDH-2.<sup>11,12</sup>

The present study has been inspired to facilitate the diagnosis prior to performing any bone marrow aspirate by estimation of the value of serum LDH in the diagnosis of megaloblastic anemia. This study has been undertaken to determine incidence of megaloblastic anemia according to age, sex and nutritional status of patients attending OPDS and wards of various departments of Sanjay Gandhi Memorial Hospital and Gandhi Memorial Hospital affiliated to Shyam Shah Medical College, Rewa, Madhya Pradesh, a tertiary care teaching hospital in Central India.

## METHODS

For the present study, the cases were selected from patients attended the OPD and admitted in Sanjay Gandhi Memorial Hospital & Gandhi Memorial Hospital, Shyam Shah Medical College, Rewa, Madhya Pradesh, a tertiary care teaching hospital in Central India. Criteria for selection of the patients were those patients presenting with anaemia.

Careful history and physical examination were done to establish the underlying cause of anaemia. Conditions known to be associated with a rise in serum LDH activity like myocardial infarction, pulmonary infarction, congestive heart failure, hepatitis, cirrhosis, extensive carcinomatosis, leukaemia etc. were excluded from the study. Following investigations were then done to classify anaemia and to establish the diagnosis of megaloblastic anaemia.

- Haemoglobin estimation by cyanmethaemoglobin method<sup>13</sup>
- PCV, RBC count and absolute values
- General blood picture
- Reticulocyte count
- Bone marrow examination
- Serum LDH estimation before and after treatment.

### *Haemoglobin estimation by cyanmethaemoglobin method (Dacie and Lewis, 2006)<sup>13</sup>*

Clinichem haemoglobin fluid stable kit was used employing cyanmethaemoglobin method and marketed by Cadila Health Care Limited Zydus Pthline Division of Ahmedabad, India.

### *Method of the test*

The test was done by colorimetric method according to the recommendation of the International Committee for standardization in hematology.

### *PCV, RBC Count and absolute values (Raphael 1976)<sup>14</sup>*

- PCV: By Wintrobe method, Apparatus: Wintrobe tube, centrifuge.
- RBC Count: By Thoma Pipette Method (Raphael 1976), Apparatus: Thoma's RBC pipette, modified Neubauer chamber.<sup>14</sup>

### *General blood picture and bone marrow aspiration and examination*

Stained by Leishman's for Giemsa's stain and then examined under light microscope to establish the type of anaemia and to confirm megaloblastic anaemia by bone marrow examination.

### *Serum LDH (lactate dehydrogenase) estimation*

LDH-UV Kinetic method was employed to the determination of lactate dehydrogenase in serum and plasma LDH Kit marketed by Reckon Diagnostic Pvt. Ltd. Baroda Lot no. 8H014 was used.

### *Principle*

LDH catalyses the oxidation of lactate to pyruvate accompanied by the simultaneous reduction of NAD to

NADH. LDH activity in serum is proportional to the increase in absorbance due to reduction of NAD.

## RESULTS

The present study was conducted with aim of assessing the incidence of megaloblastic anemia in Indian adults presenting with anemia and to evaluate the significance of serum LDH estimation as a diagnostic tool and as a prognostic indicator by its estimation subsequent to treatment and hence evaluating the response to treatment. For the above, 100 cases presenting with anemia in

Sanjay Gandhi Memorial Hospital and Gandhi Memorial Hospital were considered. Proper history and examination were done. Of the 100 cases, 50 cases (50%) of the cases were microcytic hypochromic anemia. 15 cases (15%) were normocytic normochromic anemia; 35 cases (35%) were macrocytic anemia on the basis of general blood picture and absolute values (Table 1). Of the 35 cases which were macrocytic anemia according to general blood picture and absolute values, 20 cases (57.14%) had megaloblastic bone marrow and 15 cases (42.86%) had normoblastic bone marrow (Table 2). The incidence of megaloblastic anaemia in Indian adults was 20% (Table 3).

**Table 1: Distribution of cases according to general blood picture and absolute values.**

MCV (fl)	MCH (pg)	MCHC (g/dl)	G.B.P.	No. of cases	Percentage
<80	<27	<30	Microcytic hypochromic	50	50
80 -95	27-33	30-36	Normocyte normochromic	15	15
>95	>33	30-36	Macrocytic	35	35

**Table 2: Distribution of macrocytic anemia cases on the basis of bone marrow morphology.**

Bonemarrow morphology	No. of cases	%
Megaloblastic	20	57.14
Normoblastic	15	42.86
Total	35	100%

**Table 3: Incidence of megaloblastic anemia in Indian adults.**

Type of Anaemia	No. of cases	%
Megaloblastic anemia	20	20
Non-megaloblastic anemia	80	80
Total	100	100

**Table 4: Distribution of megaloblastic anemia cases on the basis of age.**

Age (in yrs)	No. of cases	Percentage
20-30	11	55
30-40	5	25
40-50	2	10
50-60	1	5
60-70	1	5
Total	20	100

The maximum cases i.e. 55% were in 20-30 years of age followed by 25% in 30-40 years of age. Minimum cases 5% were in the age group of 50-60 years and 60-70 years (Table 4).

Of the 20 cases, 11 cases (55%) were males and 9 cases (45%) were females. Male female ratio was slightly in favour of males i.e. 1.2:1 (Table 5).

**Table 5: Distribution of megaloblastic anemia cases on the basis of sex.**

Sex	No. of cases	%
Male	11	55
Female	9	45
Total	20	100

**Table 6: Distribution of megaloblastic anemia cases on the basis of nutritional status.**

Nutritional Status	No. of cases	%
Satisfactory	4	20
Unsatisfactory	16	80
Total	20	100

Proper dietary history was taken and leading questions regarding the intake of milk, cheese, vegetables, meat etc were asked to assess whether diet included Vitamin B12 and folic acid. Those cases lacking the above were labeled as having unsatisfactory nutritional status.

**Table 7: Distribution of megaloblastic anemia cases on the basis of hemoglobin level.**

Hemoglobin level (gm/dl)	No. of cases	%
<2	2	10
2-3	5	25
3-4	4	20
4-5	3	15
5-6	3	15
6-7	2	10
>7	1	5
Total	20	100

Maximum number of cases i.e. 80% had unsatisfactory nutritional status (Table 6). Maximum number of cases i.e. 25% of the cases presented when hemoglobin level was 2-3 gm/dl followed by 20% of the cases who presented when their hemoglobin level was 3-4gm/dl. Minimum case, i.e. 5% of the cases presented when the hemoglobin was more than 7 gm/dl (Table 7). All the cases had MCV more than 95 fl while 90% of the cases had MCV more than 100 fl (Table 8).

**Table 8: Distribution of megaloblastic anemia cases on the basis of MCV.**

MCV (fl)	No. of cases	Percentage
<95	0	0
95-100	2	10
>100	18	90
Total	20	100

**Table 9: Distribution of megaloblastic anemia cases on the basis of MCH.**

MCH (Pg)	No. of cases	Percentage
<32	0	0
32 -35	2	10
35-38	2	10
38-41	13	65
41-44	3	15

Maximum number i.e. 65% of the cases, had MCH ranging from 38-41 pg, 90 %of the cases MCH more than 35 pg while only 10% of the cases had MCH of 32-35 pg, 15% of the cases had MCH as high as 41-44 pg (Table 9).

**Table 10: Distribution of megaloblastic anemia cases on the basis of MCHC.**

MCHC (gm/dl)	No. of cases	Percentage
<30	0	0
30-32	4	20
32-34	14	70
34-36	2	10
>36	0	0
Total	20	100

All 20 cases had MCHC within the normal limits (30-36) gm/dl (Table 10).

**Table 11: Distribution of megaloblastic anemia cases on the basis of presence of hyper segmented polymorphs in the peripheral blood.**

Presence	No. of cases	Percentage
Present	13	65
Absent	7	35
Total	20	100

About 5% or more of polymorphs with five lobes or at least one polymorph with 6 lobes or more were considered as hyper segmented polymorphs.

65% of the cases had hyper segmented polymorphs in the peripheral blood while 35% of the cases did not show hyper segmented polymorphs (Table 11).

**Table 12: Distribution of megaloblastic anemia cases on the basis of serum LDH levels.**

Serum LDH level (U/L)	No. of cases	%
<240	0	0
250-1000	2	10
1000-2000	3	15
2000-3000	3	15
3000-4000	6	30
4000-5000	6	30
Total	20	100

Lowest level recorded=520 U/L Highest level recorded = 4520 U/L.

Maximum number of cases (90%) of the cases had serum LDH level of more than 1000 U/L. Range of serum LDH level was 520 U/L to 4520 U/L. Thus, there was 2 to 20 fold of highest reference value (240 U/L at37 °C) rise in serum LDH level in megaloblastic anemia.

Maximum number of cases (30%) had serum LDH levels of 3000 to 4000 U/L and 4000 to 5000 U/L (Table 12).

**Table 13: Prognostic significance of serum LDH in six cases of megaloblastic anemia by assessing response to treatment after 7 days.**

Case no.	Serum LDH level before treatment	Serum LDH level after treatment
I	4520	1012
II	4398	1117
III	1488	520
IV	1791	542
V	3892	1136
VI	2454	728

Serum LDH level after treatment in all cases showed decrease in the levels though the drop was more in those cases with high initial levels in comparison to those with lower initial levels (Table 13).

## DISCUSSION

The present study was designed with the aim to study the incidence of megaloblastic anemia in Indian adult and to assess the role of serum LDH estimation as a diagnostic indicator and as a prognostic indicator by evaluating the response to treatment by its estimations subsequent to treatment.

### ***Incidence of megaloblastic anemia***

One hundred cases of anemia were studied who attended and admitted in Sanjay Gandhi Memorial Hospital & Gandhi Memorial Hospital and Associated Hospitals of Shyam Shah Medical College Rewa (M.P.).

These 100 cases were morphologically classified into macrocytic hypochromic, macrocytic and normocytic normochromic anemia on the basis of general blood picture and absolute values. In the present study, there were 50 cases (50%) of microcytic hypochromic anemia, 15 cases (15%) of normocytic normochromic anemia and 35 cases (35%) of macrocytic anemia.

Bone marrow examination of the 35 cases of macrocytic anemia was done to establish whether the erythropoiesis was megaloblastic or normoblastic. Of these (42.86%) were normoblastic macrocytic anemia. and remaining (57.14%) were megaloblastic anemia.

Thus, incidence of megaloblastic anemia in Indian adults is 20%. Most of the studies on incidence of megalablastic anemia have been done in children in our country and the incidence varies from as low as 3.1 to as high as 71.1%. Lakhotia et al, (1994) conducted a study on megaloblastic anaemia in Indian adults and found it to be 22.5%.<sup>15</sup>

### ***Age incidence***

In the present study, maximum cases (55%) were in the age group of 20 to 30 years followed by 25% cases in 30 to 40 years. The minimum cases (5%) were in the age group of 50 to 60 years and 60 to 70 years. Mukibi et al, (1992) studied 100 cases of megaloblastic anaemia. They have reported two peaks-one in third and fourth decade and other in seventh decade.<sup>16</sup>

In the present study, the higher incidence in third decade followed by fourth decade could be due to the fact that adults in this age group usually lead a very busy life and hence may ignore their diet. Moreover, due to increased popularity of junk food, proper balanced diet is usually not taken. The low incidence in seventh decade in our study could be because the symptoms of anemia in old age are usually ignored and attributed to age related weakness.

### ***Sex incidence***

In the present study, the males are found to be more affected (55%) than the females (45%). Male female ratio was slightly in favour of males i.e. 1.2.:1. Lakhotia et al, (1994) have reported that 30% of male population presenting with anaemia and 18.5% of female population with anaemia had megaloblastic anaemia and male to female ratio was in favour of males who had megaloblastic anaemia. Females suffer more from iron deficiency anaemia.<sup>15</sup>

### ***Nutritional status***

Proper history regarding dietary habits was taken from 20 cases of megaloblastic anemia. Leading questions regarding intake of cheese, egg, milk, meat, vegetables etc. were asked to assess whether their nutrition included Vitamin B12 and folic acid. Nutritional status of those cases which lacked the above was labelled as having unsatisfactory nutritional status. In the present study, maximum cases (80%) presenting with megaloblastic anemia had unsatisfactory nutritional status.

Only 20% of cases had satisfactory nutritional status. Unsatisfactory nutritional status could be due to poverty, low socio-economic status, illiteracy, ignorance and faulty food habits of majority of Indian population. According to Babior and Bunn (1994), inadequate dietary intake is more commonly associated with unbalanced diet as in alcoholics, teenagers, some infants and leads to folic acid deficiency.<sup>17</sup> Cobalamin deficiency is rare due to inadequate dietary intake and is found only in vegetarians.

### ***Hemoglobin level***

In the present study, maximum cases (25%) had hemoglobin level of 2 to 3 gm/dl followed by 20% of the cases who had hemoglobin level of 3 to 4 gm/dl. The minimum cases (5%) had hemoglobin level of more than 7 gm/dl. This late presentation may be because majority of people tend to ignore their weakness and ill health and turn up at the hospital when the weakness becomes too incapacitating. Author study justifies the citation of Lee (1993) that the degree of anemia at the time of presentation ranges from levels barely compatible with life to near normal.<sup>18</sup>

### ***Absolute values***

In the present study, all the cases had MCV of more than 95 fl of which 90% cases had MCV of more than 100 fl. MCH was more than 35 pg in 90% cases and 10% cases had MCH of 32 to 35 pg. MCHC was within normal limits (30-36 g/dl) in all the cases. This is in correspondence to the studies by Hallberg (1965) & Hall (1981).<sup>19,20</sup> Wilkinson (1949) has also reported macrocytosis with increase in MCV characteristically precedes the development of anemia.<sup>21</sup> Mukiibi et al, (1990) have reported the significance of cell indices in the diagnosis of megaloblastic anemia.<sup>16</sup> MCH of 33 to 38 pg have been reported when anemia is moderate and values as high as 56 pg when anemia is more severe (Lee, 1993).<sup>18</sup>

### ***Hyper segmented polymorphs***

About 5% or more of polymorphs with 5 lobes or at least one polymorph with 6 lobes or more were considered as hypersegmented polymorphs. In the present study, maximum number of cases (65%) showed the presence of hypersegmented polymorphs while 35% of the cases did not show hypersegmented polymorphs. This is in



correspondence with the study of Lindenbaun and Nath (1980) have reported 98% of patients with megaloblastic anemia presenting with hypersegmented polymorphs.<sup>22</sup> However, Lakhotia et al, (1994) who reported hypersegmented polymorphs in 61.1% of the case.<sup>15</sup>

### Serum LDH Level

#### Diagnostic significance

In the present study, 90% of the cases with bone marrow proven megaloblastic anemia had serum LDH level of more than 1000 U/L. The lowest level recorded was 520 U/L and the highest level recorded was 4520 U/L. The range of serum LDH values obtained in author study 520 U/L to 4520 U/L. Thus, there was 2 to 20-fold rise in the levels of serum LDH. Maximum cases (30%) had value of 3000 to 4000 U/L and 4000 to 5000 U/L.

Author findings are in correspondence with the studies of previous workers. Hess and Gehm (1955) have reported serum LDH values from 5 to 21 times the upper limit of normal in 16 cases of pernicious anemia.<sup>23</sup> Their findings were later confirmed by Zimmerman, West and Heller (1958); Heller, West and Zimmerman (1959), Gordin and Enari (1959); Levitan, Wasserman and Wroblewski (1959); Amelung (1960); Gronval (1961); Elliot and Wilkinson (1963) and Goldfarb and Papp (1963).<sup>24-31</sup>

Emerson and Wilkinson (1966) have found abnormal rise in serum LDH levels in 88% of the megaloblastic anaemia cases.<sup>32</sup> Emerson and Wilkinson (1966) have reported the serum LDH values ranging from 140 to 10000 U/L.<sup>32</sup>

Winston, Warburton and Scott (1970) have also shown serum LDH level ranging from 600-15200 U/L whether in all cases of megaloblastic anemia whether due to deficiency of Vitamin B12 or folate or mixed deficiency of both the vitamins.<sup>33</sup> Lakhotia et al, (1994) have also reported LDH levels above 1000 U/L in 77.77% of megaloblastic anaemia cases.<sup>15</sup> Jaiswal and colleagues (2000) evaluated the efficacy of total serum LDH levels and LDH isoenzyme pattern in the diagnosis of megaloblastic anemia.<sup>34</sup>

#### Prognostic significance

Prognostic significance of serum LDH was assessed by estimation of serum LDH level subsequent to treatment to determine how much the anemia is responding to treatment. For this purpose, serum or plasma was collected 7 days after the treatment is instituted. Not all patients who were diagnosed as megaloblastic anemia could be followed due to their poor compliance.

These patients stopped attending the OPD and stopped the treatment as soon as they began to improve symptomatically. Six patients could be followed because they were admitted in the wards and their serum LDH

levels estimated subsequent to treatment. All of them showed decrease in serum LDH levels.

The decrease in those cases with very high values initially was greater than in those who had lesser increase in the levels initially.

Amelung (1960) have also recorded that appropriate treatment of megaloblastic anaemia with Vitamin B12 or folic acid produces a dramatic reduction of serum LDH activity.<sup>28</sup> Emerson and Wilkinson (1966) have also reported that there is a fall in serum LDH activity by the 5th post treatment day even in patients who initially had values within or just above the normal range though the fall was slight in latter cases as compared with those in which the initial serum LDH value was markedly raised.<sup>32</sup>

Eivazi-Ziaei J, Dastgiri S, Sannt Z (2007) had reported that proportional diagnostic value was significantly higher when MPXI and serum LDH used together in diagnosis of megaloblastic anemia.<sup>35</sup> They reported that mean value of MPXI significantly decreased after treatment that was 20.4 before and -0.75 after the treatment. The same significant pattern was also observed in serum LDH the mean value of serum LDH was 4230U/L before treatment and 783U/L after treatment (p <0.001).

### CONCLUSION

Maximum number of cases presenting with megaloblastic anemia 80% had unsatisfactory nutritional status and composed of diet deficient in sources rich in Vitamin B12 and folic acid. Maximum number of cases 25% presented when hemoglobin level of 2 to 3 gm/dl. MCV was more than 100 fl in 90% cases presenting with megaloblastic anemia. MCH was more than 35 pg in 90% cases. Serum LDH level in 90% of megaloblastic anemia cases was more than 1000 U/L. The range of serum LDH in author study was 520 U/L to 4520 U/L.

The rise in serum LDH levels in megaloblastic anemia varied from 2-fold to 20-fold of the higher limit of the normal reference value. Maximum cases (30%) had serum LDH value of 3000 to 4000 and 4000 to 5000 U/L.

Six cases were followed and the serum LDH levels estimation 7 days subsequent to treatment showed that there is a decrease in the levels of those cases with very high values showed a greater drop than those cases who had lesser values previously. Thus, to conclude, megaloblastic anemia is not uncommon in Indian adults and serum LDH levels provide an important means of diagnosis. It is a non-invasive procedure, safe, and does not require any expertise.

Thus, it can be performed in the peripheral centers where bone marrow aspiration cannot be performed due to lack of expertise and proper treatment can be instituted at the earliest. Their subsequent estimation are useful aid to

assess the response to treatment and hence to assess the prognosis.

*Funding: No funding sources*

*Conflict of interest: None declared*

*Ethical approval: The study was approved by the Institutional Ethics Committee*

## REFERENCES

- Beck WS. The metabolic functions of vitamin B12. N Engl J Med. 1962;266(16):708-14.
- Chanarin I, Deacon R, Perry J, Lumb M. How vitamin B12 acts. Brit J Haematol. 1981;47(4):487-91.
- Castle WB. Current concepts of pernicious anaemia. Am J Med. 1970;48:541-48.
- Linman JW. Vit B12 and folate deficiency anaemias (Megaloblastic anaemias): Haematology, Physiologic, Pathophysiologic and clinical principles. MacMillan Publishing Co. Inc New York; 1975: 359-60.
- Baker SJ, Demoeuer EM. Nutritional anaemia, its understanding and control with special reference to the work of WHO. Am J Clin Nutr. 1979;32(2):368-408.
- Herbert V. Megaloblastic anaemias. Lab Invest. 1985;52(1):3-19.
- Lee GR. Megaloblastic and Non-Megaloblastic macrocytic anemias. Wintrobe's Clinical Hematology, Lea & Febiger Philadelphia. London. 1993;1(9):745-80.
- DeGruchy GC. The red cell and anaemia: Chemical Haematology In: Medical Practice: Firkin F, Chesterman C, Penington D, Rush B (eds) 5<sup>th</sup> Ed: Oxford University Press, Delhi; 1994: 25.
- Jandl JH. Megaloblastic Anemias. Blod Test-Bood of Haematolgy, 2<sup>nd</sup> Ed. Little Brown and Company, Boston, New York, Toronto London; 1996: 251-280.
- Libnoch JA, Yakulis VJ and Heller P. Lactate dehydrogenase in megaloblastic none marrow. Am J Clin Patho. 1966;45(3):302-5.
- Amelung D. Serum enzyme determinations in pernicious anemia [Article in German]. Dtsch Med Wochenschr. 1960 Sep 9;85:1629-32.
- Wroblewski F, Gregory KF. Lactic dehydrogenase isoenzymes and their distribution in normal tissues and plasma and in disease states. Ann N Y Acad Sci. 1961;94:912-32.
- Dacie JV, Lewis SM. Basic hematological techniques In: Practical Haematology, 10<sup>th</sup> Ed. Churchill Livingstone, Edinburgh, London, Melbourne and New York; 2006: 26-30.
- Raphael SS. Basic Hematologic Techniques In: Lynch's Medical Laboratory Technology, 3<sup>rd</sup> Ed, Wb Saunder's Company, Philedelphia, London, Toronto; 1976: 1073-1129.
- Lakhotia M, Shah PKD, Balani V, Gupta A, Saxena A, Sinha HV. Incidence of megaloblastic anemia in Indian adults and the role of serum LDH as a diagnostic tool Indian J Haematol Blood Trans. 1994;12:113-5.
- Mukiibi JM, Makumbi FA, Gwanzura C. Megaloblastic anemia in Zimbabwe: Spectrum of clinical and hematological manifestations. East Afr Med J. 1992;69(2):83-7.
- Babior BM, Bunn HFB. Megaloblastic anemias. In: Isselbacher KJ, Braunwald E, Wilson JD, Martin JB, Fauci AS, Kasper DL, eds. Harrison's Principles of Internal Medicine, 13<sup>th</sup> Ed. McGraw Hill Inc. New York, St. Louis, San Francisco, Tokyo, Toronto; 1994: 1726.
- Lee GR, Foerster J, Lukens J, Paraskevas F, Greer JP, Rodgers GM. Shirley Parker Levine-Megaloblastic anaemia. In: Lipincott Williams and Wilkins,eds. Wintrobe's Clin Haematol. Vol. 2, 10<sup>th</sup> ed. Philadelphia;1999:1579-82.
- Hallberg L. Blood volume, hemolysis and regeneration of blood in pernicious anemia; studies based on the endogenous formation of carbon monoxide and determinations of the total amount of hemoglobin. Scand J Clin Lab Invest. 1955;7(16):1-127.
- Hall CA. Vitamin B12 deficiency and early rise in mean corpuscular volume. JAMA. 1981 20;245(11):1144-6.
- Wilkinson J. Megalocytic anaemias. Lancet. 1949;253(6548):336-40.
- Lindenbaum J, Nath BJ. Megaloblastic anaemia and neutrophil hypersegmentation. Brit J Haem. 1980;44(3);511.
- Hess B, Gehm E. Lactic acid dehydrogenase in the human blood. Klin Wochenschr. 1955;33(3-4):91-3.
- Zimmerman HJ, West M and Heller P. Serum enzymes in disease-II lactic dehydrogenase and glutamic oxaloacetic transaminase in anemia. AMA Arch Int Med. 1958;102(1):115-123.
- Heller H, Hollmann W, Knapp A. Iron deficiency anemia in young women. Z Gesamte Inn Med. 1959;14:1105-8.
- Gordin R, Enari TM. Lactic acid dehydrogenase in megaloblastic anaemia. The effect of corticosteroid therapy. Acta Haematol. 1959;21(6):360-5.
- Levitan R, Wasserman LR, Wroblewski E. Clinical significance of serum enzyme alteration in hematologic diseases. Clin Res. 1959;7:217.
- Amelung D. Serum ferment bestimmungen bei pernizioser Anamic Serum enzyme determinations in pernicious anemia. Dtsch Med Wochenschr. 1960 Sep 9;85:1629-32.
- Gronvall C. On the serum activity of lactic acid dehydrogenase and phosphohexose isomerase in pernicious and hemolytic anemias. Scand J Clin Lab Invest. 1961;13(1):29-36.
- Elliot BA, Wilkinson JH. The serum alpha-hydroxybutyrate dehydrogenase in diseases other than myocardial infarction. Clin Sci. 1963;24:343-55.
- Goldfarb TG, Papp BJ. Excessively high levels of lactic acid dehydrogenase activity in pernicious anemia. Am J Med. 1963;34(4):578-82.
- Emerson PM, Wilkinson JH. Lactate dehydrogenase in the diagnosis and assessment of response to

- treatment of megaloblastic anemia. *Brit J Haemat*. 1966;12(6):678-88.
33. Winston RM, Warburton FG, Scott A. Enzymatic diagnosis of megaloblastic anemia. *Brit J Haematol*. 1970;19(5):587.
34. Jaswal TS, Mehta HC, Gupta V, Singh M, Singh S. Serum lactate dehydrogenase in diagnosis of megaloblastic anaemia. *Indian J Pathol Microbiol*. 2000;43(3):325-9.
35. Eivazi-Ziaei J, Dastgiri S, Sanaat Z. Estimation of the diagnostic value of myeloperoxidase index and

lactate dehydrogenase in megaloblastic anaemia. *J Clin Diagn Res*. 2007;1(5):380-4.

**Cite this article as:** Chakravarty N, Santhikiran D, Brahma A, Singh UR, Kol PC, Sen S. Study of serum lactate dehydrogenase level as diagnostic and prognostic indicator of megaloblastic anemia. *Int J Adv Med* 2019;6:1199-206.